

B U L L E T I N

DE LA SOCIÉTÉ DES SCIENCES
ET DES LETTRES DE ŁÓDŹ

SÉRIE:
RECHERCHES SUR LES DÉFORMATIONS

Volume LXVIII, no. 2

B U L L E T I N

DE LA SOCIÉTÉ DES SCIENCES ET DES LETTRES DE ŁÓDŹ

SÉRIE: RECHERCHES SUR LES DÉFORMATIONS

Volume LXVIII, no. 2

Rédacteur en chef et de la Série: JULIAN ŁAWRYNOWICZ

Comité de Rédaction de la Série

R. ABLAMOWICZ (Cookeville, TN), Aleksandra BASZCZYŃSKA (Łódź)
O. MARTIO (Helsinki), B. SENDOV (Sofia), D. SHOIKHET (Karmiel)
O. SUZUKI (Tokyo), M. VACCARO (Salerno), E. VESENTINI (Torino)
L. WOJTCZAK (Łódź), Ilona ZASADA (Łódź)

Secrétaire de la Série:
EWELINA FRĄTCZAK



ŁÓDŹ 2018

LÓDZKIE TOWARZYSTWO NAUKOWE

PL-90-505 Łódź, ul. M. Curie-Skłodowskiej 11

tel. (42) 66-55-459, fax (42) 66 55 464

sprzedaż wydawnictw: tel. (42) 66 55 448, <http://sklep.ltn.lodz.pl>

e-mail: biuro@ltn.lodz.pl; <http://www.ltn.lodz.pl>

REDAKCJA NACZELNA WYDAWNICTW

LÓDZKIEGO TOWARZYSTWA NAUKOWEGO

Krystyna Czyżewska, Wanda M. Krajewska (redaktor naczelny)

Edward Karasiński, Henryk Piekarski, Jan Szymczak



Ministerstwo Nauki
i Szkolnictwa Wyższego

”Bulletin de la Société des sciences et des lettres de Łódź, Série: Recherches sur les déformations” – zadanie finansowane w ramach umowy 597/P-DUN/2017 ze środków Ministra Nauki i Szkolnictwa Wyższego przeznaczonych na działalność upowszechniającą naukę

”Bulletin de la Société des sciences et des lettres de Łódź, Série: Recherches sur les déformations” – task financed under the agreement 597/P-DUN/2017 by the Minister of Science and Higher Education from the funds for science dissemination

© Copyright by Łódzkie Towarzystwo Naukowe, 2018

PL ISSN 0459-6854; e-ISSN 2450-9329

DOI: 10.26485/0459-6854; DOI: 10.26485/0459-6854/2018/68.2

Wydanie 1

Nakład 80 egz.

Przygotowanie czasopisma do digitalizacji: Justyna Walewska

The journal is indexed in: *Index Copernicus*, *PBN/POL-Index*, *EBSCOhost*, *Zentralblatt MATH*, *MathSciNet (Mathematical Reviews)* and *Referativny Zhurnal*

Druk czasopisma sfinansowany przez Uniwersytet Łódzki

TABLE DES MATIÈRES

J. Ławrynowicz and J. J. Trokhimchuk , Our friend Professor Yurii B. Zelinskii	11–12
1. A. V. Pokrovskii and O. D. Trofymenko , Mean value theorems for solutions of linear partial differential equations with constant coefficients	13–24
2. A. V. Plaksa and V. S. Shpakivskyi , Integral theorems for monogenic functions in an infinite-dimensional space with a commutative multiplication	25–36
3. A. Bakhtin , L. Vyhivska , and I. Denega , Inequality for the inner radii of symmetric non-overlapping domains	37–44
4. A. S. Serdyuk and T. A. Stepanyuk , Lebesgue-type inequalities for the Fourier sums on classes of generalized Poisson integrals	45–52
5. B. A. Klishchuk , Weakly m -convex sets and the shadow problem	53–60
6. S. Yu. Favorov and N. P. Girya , Convergence of Dirichlet series on a finite-dimensional space	61–70
7. S. V. Gryshchuk , On some cases of plane orthotropy	71–76
8. T. M. Osipchuk , On a system of balls with equal radii generating shadow at a point	77–84
9. M. V. Stefanchuk , A generalization of the concept of convexity in a hyper-complex space	85–94
10. A. Futa and D. Partyka , The Schwarz-type inequality for harmonic functions of the unit disc satisfying a sectorial condition	95–108
11. O. Suzuki , J. Ławrynowicz , M. Nowak-Kępczyk and M. Zubert , Some geometrical aspects of binary, ternary, quaternary, quinary, and senary structures in physics	109–122

B U L L E T I N

DE LA SOCIÉTÉ DES SCIENCES ET DES LETTRES DE LÓDŹ

2018

Vol. LXVIII

Recherches sur les déformations

no. 2

pp. 71–76

*Dedicated to the memory of
Professor Yurii B. Zelinskii**Serhii V. Gryshchuk*

ON SOME CASES OF PLANE ORTHOTROPY

Summary

There are considered some cases of plane orthotropy in the absence of body forces. Then every function from a pair-solution of the equilibrium system of equations with respect to displacements satisfies the elliptic fourth-order equation of the type:

$$\left(\alpha_1 \frac{\partial^4}{\partial x^4} + \alpha_2 \frac{\partial^4}{\partial x^2 \partial y^2} + \frac{\partial^4}{\partial y^4} \right) w(x, y) = 0,$$

with certain real $\alpha_k \neq 0$, $k = 1, 2$.

Keywords and phrases: the generalized Hook's law, a plane orthotropy, the equilibrium system

1. Introduction

As well-known (cf., e.g., [1, 2, 3]), in the case of isotropic plane deformations with the absence of body forces a function (displacement) u or v from a pair-solution $(u(x, y), v(x, y))$ of the equilibrium system of equations in displacements

$$\begin{cases} (\lambda + \mu) \left(\frac{\partial^2 u(x, y)}{\partial x^2} + \frac{\partial^2 v(x, y)}{\partial x \partial y} \right) + \mu \Delta_2 u(x, y) = 0, \\ (\lambda + \mu) \left(\frac{\partial^2 u(x, y)}{\partial x \partial y} + \frac{\partial^2 v(x, y)}{\partial y^2} \right) + \mu \Delta_2 v(x, y) = 0 \forall (x, y) \in D, \end{cases} \quad (1)$$

as well as the stress Airy's function, satisfies the biharmonic equation: $(\Delta_2)^2 w(x, y) = 0$, where $\Delta_2 := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ is the 2-D Laplasian, D is a domain of the Cartesian plane xOy , λ and μ are the Lamé constants.

Similar results for anisotropic solid body are not well-known. One of the reason of this fact is a difficulty (due to a variety of coefficients) of the generalized Hooke's law expressing strains via stresses in a linear form.

The aim of this paper is to prove analogous (to the isotropic case) statements for some cases of an elastic anisotropic homogeneous plane solid body — a plane orthotropic body, or briefly, a plane orthotropy. We will restrict our attention on some simple but interesting cases of orthotropy.

2. Notations and preliminaries

Let $\mathbb{R}^{3 \times 3}$ be a set of all real 3×3 matrices, $A \in \mathbb{R}^{3 \times 3}$, $\det A$ is a determinant of A . If $\det A \neq 0$ then there exists the inverse matrix $B = A^{-1}$ such that $AB = BA = 1$, where 1 is the unity matrix. By $\mathbb{R}_+^{3 \times 3}$ we define all matrices of $\mathbb{R}^{3 \times 3}$ which are symmetric and positive defined. A symbol $\overleftarrow{\vartheta}$ defines a vector-column having three real coordinates ϑ_k , $k = 1, 2, 3$.

Let a model of an elastic anisotropic medium occupied a domain D of the Cartesian plane xOy be a homogeneous (cf., e.g., [4, p. 25]) plane orthotropic (cf., e.g., [4, p. 35]) body.

Let $\overleftarrow{\varepsilon}$ has coordinates equal to strains (cf., e.g., [4, p. 18]):

$$\varepsilon_1 := \varepsilon_x, \varepsilon_2 := \varepsilon_y, \varepsilon_3 = \gamma_{xy}.$$

Let $\overleftarrow{\sigma}$ has coordinates equal to stresses (cf., e.g., [4, p. 16]):

$$\sigma_1 := \sigma_x, \sigma_2 := \sigma_y, \sigma_3 := \tau_{xy}.$$

The generalized Hooke's law for our model has two equivalent forms (cf., e.g., [4, § 3], [5, § 4.1.3]):

$$\overleftarrow{\varepsilon} = A \overleftarrow{\sigma}, \overleftarrow{\sigma} = A^{-1} \overleftarrow{\varepsilon}, \quad (2)$$

with $A \in \mathbb{R}_+^{3 \times 3}$ of the form

$$A = \begin{pmatrix} a_{11} & a_{12} & 0 \\ a_{12} & a_{22} & 0 \\ 0 & 0 & a_{66} \end{pmatrix}, \quad (3)$$

where

$$a_{11} > 0, a_{11}a_{22} - (a_{12})^2 > 0, a_{66} > 0. \quad (4)$$

Unequalities (4) follows from the Sylvester's criterion of positive definiteness of the matrix (3).

A numbers a_{ij} and A_{ij} , $k \leq m$, $k, m = 1, 2, 6$, are called *elastic constants* ([4, p. 27]). They are constants in D due to the homogeneity of the solid body.

Consider notations for elements of A^{-1} :

$$A^{-1} =: \begin{pmatrix} A_{11} & A_{12} & 0 \\ A_{12} & A_{22} & 0 \\ 0 & 0 & A_{66} \end{pmatrix}, \quad (5)$$

where A_{km} satisfy (4) with $a_{km} := A_{km}$, $k \leq m$, $k, m = 1, 2, 6$.

A *stress function* ([6, p. 21] with $\bar{U} \equiv 0$) is a function w satisfying relations:

$$\begin{aligned} \sigma_x(x_0, y_0) &= \frac{\partial^2 w}{\partial y^2}(x_0, y_0), \quad \sigma_y(x_0, y_0) = \frac{\partial^2 w}{\partial x^2}(x_0, y_0), \\ \tau_{xy}(x_0, y_0) &= -\frac{\partial^2 w}{\partial x \partial y}(x_0, y_0) \quad \forall (x_0, y_0) \in D. \end{aligned}$$

In the absence of body forces, the stress function $w(x, y)$ satisfies the elliptic fourth-order equation (“the stress equation”, cf., e.g., [6, p. 27] with $a_{16} = a_{26} = 0$):

$$\left(a_{22} \frac{\partial^4}{\partial x^2} + (2a_{12} + a_{66}) \frac{\partial^4}{\partial x^2 \partial y^2} + a_{11} \frac{\partial^4}{\partial y^4} \right) w(x, y) = 0. \quad (6)$$

The equilibrium system of equations with respect to the displacement vector $(u(x, y), v(x, y))$ has a form (cf., e.g., [4, p. 75]):

$$\begin{cases} \left(A_{11} \frac{\partial^2}{\partial x^2} + A_{66} \frac{\partial^2}{\partial y^2} \right) u(x, y) + (A_{12} + A_{66}) \frac{\partial^2 v(x, y)}{\partial x \partial y} = 0, \\ \left(A_{66} \frac{\partial^2}{\partial x^2} + A_{22} \frac{\partial^2}{\partial y^2} \right) v(x, y) + (A_{12} + A_{66}) \frac{\partial^2 u(x, y)}{\partial x \partial y} = 0, \end{cases} \quad (7)$$

where all $(x, y) \in D$; A_{km} , $k \leq m$, $k, m = 1, 2, 6$, are defined in (5).

3. Cases of orthotropy and solutions of their equilibrium systems and stress equation

Consider the following equation (particular case of (6)):

$$\begin{aligned} & l_{0,p} w(x, y) \equiv \\ & \equiv \left((2p - 1) \frac{\partial^4}{\partial x^4} + 2p \frac{\partial^4}{\partial x^2 \partial y^2} + \frac{\partial^4}{\partial y^4} \right) w(x, y) = 0 \quad \forall (x, y) \in D, \end{aligned} \quad (8)$$

where $p \neq 1$ is a real parameter.

Consider an orthotropy with

$$a_{11} = a_{12} = 1, \quad a_{22} = 2p - 1, \quad a_{16} = a_{26} = 0, \quad a_{66} = 2(p - 1). \quad (9)$$

Then the equation (8) is a stress equation. It is easy to check that the matrix (3) is positive defened only for $p > 1$. So a case $p < 1$ has no elastic meaning and we are to investigate a case $p > 1$. Calculating the inverse matrix A^{-1} we find:

$$A_{11} = \frac{2p - 1}{2(p - 1)}, \quad A_{12} = -\frac{1}{2(p - 1)}, \quad A_{22} = A_{66} = -A_{12}. \quad (10)$$

Since $A_{12} + A_{66} = 0$ a system (7) takes a form

$$\begin{cases} \frac{1}{2(p-1)} l_{1,p} u(x, y) = 0, \\ \frac{1}{2(p-1)} \Delta_2 v(x, y) = 0 \forall (x, y) \in D, \end{cases} \quad (11)$$

where $l_{1,p} := (2p-1) \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$.

Taking into account that the operator (8) can be factorised in the form:

$$l_{1,p} = l_{1,p} \circ \Delta_2 = \Delta_2 \circ l_{1,p}$$

($l_1 \circ l_2$ is a symbol of composition of operators l_1 and l_2), we see that if a pair (u, v) is a solution of (11) then $w := u$ or $w := v$ is a solution of the equation (8). So we proved the following theorem.

Theorem 1. *Let $p > 1$, an orthotropy is defined by (2), (9). Then every displacement-function from a pair of solution of the equilibrium system (11) satisfies the equation (8).*

Now consider another cases of orthotropy for which an equilibrium equation splits onto two equations containing except of operators of the type $l_{1,p}$ an extra term-operator $\frac{\partial^2}{\partial x \partial y}$ acted to another unknown function and has a non-zero coefficient.

Let p be an arbitrary fixed number: $0 < p < 1$.

Take into consideration the plane orthotropy:

$$a_{11} = a_{22} = 1, a_{16} = a_{26} = 0, a_{66} = 2(p - a_{12}), -1 < a_{12} < p. \quad (12)$$

An a_{12} belongs to such measures due to the positiveness of the matrix (3). Therefore, we have:

$$A_{11} = A_{22} = \frac{1}{1 - a_{12}^2}, A_{21} = A_{12} = -\frac{a_{12}}{1 - a_{12}^2}, A_{66} = \frac{1}{2(p - a_{12})}.$$

The equilibrium system (7) gets a form:

$$\begin{cases} \frac{1}{1 - a_{12}^2} \frac{\partial^2}{\partial x^2} u(x, y) + \frac{1}{2(p - a_{12})} \frac{\partial^2}{\partial y^2} u(x, y) + \\ + \left(-\frac{a_{12}}{1 - a_{12}^2} + \frac{1}{2(p - a_{12})} \right) \frac{\partial^2 v(x, y)}{\partial x \partial y} = 0, \\ \frac{1}{2(p - a_{12})} \frac{\partial^2}{\partial x^2} u(x, y) + \frac{1}{1 - a_{12}^2} \frac{\partial^2}{\partial y^2} v(x, y) + \\ + \left(-\frac{a_{12}}{1 - a_{12}^2} + \frac{1}{2(p - a_{12})} \right) \frac{\partial^2 u(x, y)}{\partial x \partial y} = 0, \end{cases} \quad (13)$$

where all $(x, y) \in D$.

Consider the following (“stress”) equation:

$$\begin{aligned} & l_{2,p} w(x, y) \equiv \\ & \equiv \left(\frac{\partial^4}{\partial x^4} + 2p \frac{\partial^4}{\partial x^2 \partial y^2} + \frac{\partial^4}{\partial y^4} \right) w(x, y) = 0 \forall (x, y) \in D. \end{aligned} \quad (14)$$

The system (13) is equivalent to the following system:

$$\begin{cases} B_{11} \frac{\partial^2 u(x,y)}{\partial x^2} + B_{12} \frac{\partial^2 u(x,y)}{\partial y^2} + \frac{\partial^2 v(x,y)}{\partial x \partial y} = 0, \\ B_{21} \frac{\partial^2 v(x,y)}{\partial x^2} + B_{22} \frac{\partial^2 v(x,y)}{\partial y^2} + \frac{\partial^2 u(x,y)}{\partial x \partial y} = 0 \forall (x,y) \in D. \end{cases} \quad (15)$$

where

$$B_{11} = B_{22} := \frac{2(p - a_{12})}{(a_{12} - p)^2 + 1 - p^2},$$

$$B_{12} = B_{21} := \frac{1 - (a_{12})^2}{(a_{12} - p)^2 + 1 - p^2}.$$

Theorem 2. *Let $0 < p < 1$, an orthotropy is defined by (2), (12). Then every displacement-function from a pair of solution of the equilibrium system (15) satisfies the equation (14).*

Proof. Acting by the differential operator $\frac{\partial^2}{\partial x \partial y}$ on the second equation of (15) and substituting to the obtained equation an expression of $\frac{\partial^2 v}{\partial x \partial y}$, we arrive at the equation:

$$\frac{\partial^4 u(x,y)}{\partial x^4} + C_2 \frac{\partial^4 u(x,y)}{\partial x^2 \partial y^2} + C_3 \frac{\partial^4 u(x,y)}{\partial y^4} \forall (x,y) \in D, \quad (16)$$

where

$$C_3 = \frac{B_{22} B_{12}}{B_{11} B_{21}} \equiv 1, \quad C_2 := \frac{B_{11} B_{22} + B_{12} B_{21} - 1}{B_{11} B_{21}}.$$

So, to prove Theorem we need to check the equality $C_2 = 2p$. In terms of p and a_{12} the relation $C_2 = 2p$ can be rewritten in the form:

$$\alpha^2 + \beta^2 + 2p\alpha\beta = (\alpha + a_{12}\beta)^2,$$

where $\alpha := 1 - a_{12}^2$, $\beta := 2(a_{12} - p)$. By doing simple algebraic transformation, the last one is equivalent to the relation

$$1 - a_{12}^2 = 2(a_{12} - p) \frac{\alpha}{\beta},$$

which with use of the definitions of α and β is an identity. So, we proved that if (u, v) is a solution of (15) then v satisfies the equation (14).

A similar statement for v can be proved analogously. The theorem is proved. \square

Acknowledgements

The research is partially supported by the Polish-Ukrainian grant No. 39/2014 *Topological-analytical methods in complex and hypercomplex analysis* of the Polish Academy of Sciences and the National Academy of Sciences of Ukraine and by the Fundamental Research Programme funded by the Ministry of Education and Science of Ukraine (project No. 0116U001528).

References

- [1] S. G. Mikhlin, *The plane problem of the theory of elasticity*, Trans. Inst. of seismology, Acad. Sci. USSR, No. 65, Acad. Sci. USSR Publ. House, Moscow–Leningrad, 1935 (in Russian).
- [2] N. I. Muskhelishvili, *Some basic problems of the mathematical theory of elasticity. Fundamental equations, plane theory of elasticity, torsion and bending*, Springer Science+Business Media Dordrecht, Originally published by Noordhoff International Publishing in 1977, 1977.
- [3] J. Lu, *Complex Variable Methods in Plane Elasticity /Series in Pure Mathematics. Vol. 22/*, World Scientific, Singapore, 1995.
- [4] S. G. Lekhnitskii, *Theory of Elasticity of an Anisotropic Body* (Engl. transl. from the revised 1977 Russian edition), “MIR” Publ., 1981.
- [5] P. P. Teodorescu, *Treatise on Classical Elasticity Theory and Related Problems /Mathematical and Analytical Techniques with Applications to Engineering/* (Ed. A. Jeffrey), Springer, Berlin, 2010.
- [6] S. G. Lekhnitskii, *Anisotropic Plates* (Engl. transl. from the 2nd Russ. edition by S. W. Tsai, T. Cheron), Gordon and Breach, New York, 1984.

Institute of Mathematics
National Academy of Sciences of Ukraine
Tereshchenkivska str. 3, UA-01004, Kyiv
Ukraine
E-mail: serhii.gryshchuk@gmail.com

Presented by Julian Ławrynowicz at the Session of the Mathematical-Physical Commission of the Łódź Society of Sciences and Arts on June 27, 2018.

O PEWNYCH PRZYPADKACH PŁASKIEJ ORTHOTROPII

S t r e s z c z e n i e

Rozpatrywane są pewne przypadki płaskiej orthotropii przy założeniu braku oddziaływania sił ciała. Wówczas każda funkcja z pary rozwiązań układu równowagi równań ze względu na przemieszczenia spełnia równanie eliptyczne czwartego rzędu typu:

$$\left(\alpha_1 \frac{\partial^4}{\partial x^4} + \alpha_2 \frac{\partial^4}{\partial x^2 \partial y^2} + \frac{\partial^4}{\partial y^4} \right) w(x, y) = 0,$$

z pewnymi rzeczywistymi stałymi $\alpha_k \neq 0$, $k = 1, 2$.

Słowa kluczowe: uogólnione prawo Hooke’a, orthotropia płaska, układ równowagi