

# Generalized integral theorems for the quaternionic $G$ -monogenic mappings

TETYANA S. KUZMENKO AND VITALII S. SHPAKIVSKYI

(Presented by V. Ya. Gutlyanskii)

**Abstract.** For  $G$ -monogenic mappings taking values in the algebra of complex quaternions we generalize some analogues of classical integral theorems of the holomorphic function theory of a complex variable (the surface and the curvilinear Cauchy integral theorems).

**2010 MSC.** 30G35, 57R35.

**Key words and phrases.** The algebra of complex quaternions,  $G$ -monogenic mappings, the Cauchy integral theorems.

## 1. Introduction

The Cauchy integral theorems for analytic functions of the complex variable are fundamental results of the classical complex analysis. Analogues of these results are also important tools in the quaternionic analysis.

In the papers [1–3], some analogues of classical integral theorems for  $G$ -monogenic mappings taking values in the algebra of complex quaternions were established. Namely, in the paper [1] the Stokes formula, a curvilinear analogue of the Cauchy integral theorem in the case where a curve of integration lies in a domain of  $G$ -monogeneity, the Cauchy integral formula, the Gauss–Ostrogradsky formula and the surface Cauchy integral theorem were proved. The analogues of the Cauchy integral theorems are of the form

$$\int_{\Gamma} \widehat{\Phi} \sigma = 0, \quad \int_{\Gamma} \sigma \Phi = 0, \quad (1.1)$$

---

Received 24.10.2016

*This research is partially supported by Grant of Ministry of Education and Science of Ukraine (Project No. 0116U001528).*

where  $\Gamma$  is a closed surface (or a closed curve),  $\sigma$  is a special differential form, and  $\widehat{\Phi}$ ,  $\Phi$  are left- $G$ -monogenic mapping and right- $G$ -monogenic mapping, respectively.

In the paper [2], the formulae (1.1) was proved in the case where a curve of integration lies on the boundary of a domain of  $G$ -monogeneity. In the paper [3], the analogue of the Morera theorem was established.

In the present paper we generalize analogues of the surface and curvilinear Cauchy integral theorems for  $G$ -monogenic mappings to “two sides” integrals. Namely, the equality

$$\int_{\Gamma} \widehat{\Phi} \sigma \Phi = 0 \tag{1.2}$$

will be proved under some assumptions. In the papers [4] and [5] the formula of the type (1.2) was proved for another class of quaternionic differentiable functions.

## 2. $G$ -monogenic mappings in the algebra of complex quaternions

Let  $\mathbb{H}(\mathbb{C})$  be the quaternion algebra over the field of complex numbers  $\mathbb{C}$ , whose basis consists of the unit 1 of the algebra and of the elements  $I, J, K$  satisfying the multiplication rules:

$$I^2 = J^2 = K^2 = -1,$$

$$IJ = -JI = K, \quad JK = -KJ = I, \quad KI = -IK = J.$$

In the algebra  $\mathbb{H}(\mathbb{C})$  there exists another basis  $\{e_1, e_2, e_3, e_4\}$  such that multiplication table in this basis can be represented as

$\cdot$	$e_1$	$e_2$	$e_3$	$e_4$
$e_1$	$e_1$	$0$	$e_3$	$0$
$e_2$	$0$	$e_2$	$0$	$e_4$
$e_3$	$0$	$e_3$	$0$	$e_1$
$e_4$	$e_4$	$0$	$e_2$	$0$

The unit of the algebra can be decomposed as  $1 = e_1 + e_2$ .

Let us consider the vectors

$$i_1 = e_1 + e_2, \quad i_2 = a_1 e_1 + a_2 e_2, \quad i_3 = b_1 e_1 + b_2 e_2, \tag{2.1}$$

where  $a_k, b_k \in \mathbb{C}$ ,  $k = 1, 2$ , which are linearly independent over the field of real numbers  $\mathbb{R}$ . It means that the equality  $\alpha_1 i_1 + \alpha_2 i_2 + \alpha_3 i_3 = 0$  for  $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$  holds if and only if  $\alpha_1 = \alpha_2 = \alpha_3 = 0$ .

In the algebra  $\mathbb{H}(\mathbb{C})$  we consider the linear span

$$E_3 := \{\zeta = xi_1 + yi_2 + zi_3 : x, y, z \in \mathbb{R}\}$$

generated by the vectors  $i_1, i_2, i_3$  over the field  $\mathbb{R}$ .

A set  $S \subset \mathbb{R}^3$  is associated with the set  $S_\zeta := \{\zeta = xi_1 + yi_2 + zi_3 : (x, y, z) \in S\}$  in  $E_3$ . We understand topological properties of the set  $S_\zeta$  in  $E_3$  as the same topological properties of the set  $S$  in  $\mathbb{R}^3$ .

In the paper [6] we introduced a new class of quaternionic mappings, so-called,  $G$ -monogenic mappings.

We say that a continuous mapping  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  (or  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$ ) is *right- $G$ -monogenic* (or *left- $G$ -monogenic*) in a domain  $\Omega_\zeta \subset E_3$ , if  $\Phi$  (or  $\widehat{\Phi}$ ) is differentiable in the sense of the Gâteaux at every point of  $\Omega_\zeta$ , i. e. for every  $\zeta \in \Omega_\zeta$  there exists an element  $\Phi'(\zeta) \in \mathbb{H}(\mathbb{C})$  (or  $\widehat{\Phi}'(\zeta) \in \mathbb{H}(\mathbb{C})$ ) such that

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0+0} \left( \Phi(\zeta + \varepsilon h) - \Phi(\zeta) \right) \varepsilon^{-1} = h\Phi'(\zeta) \quad \forall h \in E_3 \\ &\left( \text{or } \lim_{\varepsilon \rightarrow 0+0} \left( \widehat{\Phi}(\zeta + \varepsilon h) - \widehat{\Phi}(\zeta) \right) \varepsilon^{-1} = \widehat{\Phi}'(\zeta)h \quad \forall h \in E_3 \right). \end{aligned}$$

Consider the decomposition of the mapping  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  with respect to the basis  $\{e_1, e_2, e_3, e_4\}$ :

$$\Phi(\zeta) = \sum_{k=1}^4 U_k(x, y, z)e_k.$$

In the case where functions  $U_k : \Omega \rightarrow \mathbb{C}$  are  $\mathbb{R}$ -differentiable in  $\Omega$ , i. e. for every  $(x, y, z) \in \Omega$

$$\begin{aligned} &U_k(x + \Delta x, y + \Delta y, z + \Delta z) - U_k(x, y, z) \\ &= \frac{\partial U_k}{\partial x} \Delta x + \frac{\partial U_k}{\partial y} \Delta y + \frac{\partial U_k}{\partial z} \Delta z + o\left(\sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2}\right), \\ &(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \rightarrow 0, \end{aligned}$$

the mapping  $\Phi$  is *right- $G$ -monogenic* and  $\widehat{\Phi}$  is *left- $G$ -monogenic* in the domain  $\Omega_\zeta$  if and only if the following analogues of the Cauchy – Riemann conditions are satisfied in  $\Omega_\zeta$ :

$$\frac{\partial \Phi}{\partial y} = i_2 \frac{\partial \Phi}{\partial x}, \quad \frac{\partial \Phi}{\partial z} = i_3 \frac{\partial \Phi}{\partial x} \tag{2.2}$$

and

$$\frac{\partial \widehat{\Phi}}{\partial y} = \frac{\partial \widehat{\Phi}}{\partial x} i_2, \quad \frac{\partial \widehat{\Phi}}{\partial z} = \frac{\partial \widehat{\Phi}}{\partial x} i_3. \tag{2.3}$$

### 3. Cauchy integral theorem for a surface integral

Let  $\Omega_\zeta$  be a bounded domain in  $E_3$ . For a continuous mapping  $\varphi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  of the form

$$\varphi(\zeta) = \sum_{k=1}^4 U_k(x, y, z)e_k + i \sum_{k=1}^4 V_k(x, y, z)e_k,$$

where  $(x, y, z) \in \Omega$  and  $U_k : \Omega \rightarrow \mathbb{R}$ ,  $V_k : \Omega \rightarrow \mathbb{R}$ , we define a *volume integral* by the equality

$$\int_{\Omega_\zeta} \varphi(\zeta) dx dy dz := \sum_{k=1}^4 e_k \int_{\Omega} U_k(x, y, z) dx dy dz$$

$$+ i \sum_{k=1}^4 e_k \int_{\Omega} V_k(x, y, z) dx dy dz.$$

Let  $\Sigma_\zeta$  be a piece-smooth surface in  $E_3$ . For continuous mappings  $\varphi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and  $\psi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  of the forms

$$\varphi(\zeta) = \sum_{k=1}^4 U_k(x, y, z)e_k + i \sum_{k=1}^4 V_k(x, y, z)e_k, \quad (3.1)$$

$$\psi(\zeta) = \sum_{m=1}^4 P_m(x, y, z)e_m + i \sum_{m=1}^4 Q_m(x, y, z)e_m, \quad (3.2)$$

where  $(x, y, z) \in \Sigma$ ,  $U_k : \Sigma \rightarrow \mathbb{R}$ ,  $V_k : \Sigma \rightarrow \mathbb{R}$  and  $P_m : \Sigma \rightarrow \mathbb{R}$ ,  $Q_m : \Sigma \rightarrow \mathbb{R}$ , we define a *surface integral* on a piece-smooth surface  $\Sigma_\zeta$  with the differential form

$$\sigma := dydz + dzdx i_2 + dx dy i_3$$

by the equality

$$\begin{aligned}
 \int_{\Sigma_\zeta} \varphi(\zeta) \sigma \psi(\zeta) &:= \sum_{k,m=1}^4 e_k e_m \int_{\Sigma} (U_k P_m - V_k Q_m) dydz \\
 &+ \sum_{k,m=1}^4 e_k i_2 e_m \int_{\Sigma} (U_k P_m - V_k Q_m) dzdx \\
 &+ \sum_{k,m=1}^4 e_k i_3 e_m \int_{\Sigma} (U_k P_m - V_k Q_m) dx dy \\
 &+ i \sum_{k,m=1}^4 e_k e_m \int_{\Sigma} (V_k P_m + U_k Q_m) dydz \\
 &+ i \sum_{k,m=1}^4 e_k i_2 e_m \int_{\Sigma} (V_k P_m + U_k Q_m) dzdx \\
 &+ i \sum_{k,m=1}^4 e_k i_3 e_m \int_{\Sigma} (V_k P_m + U_k Q_m) dx dy.
 \end{aligned}$$

If a domain  $\Omega \subset \mathbb{R}^3$  has a closed piece-smooth boundary  $\partial\Omega$  and mappings  $\varphi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and  $\psi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  are continuous together with partial derivatives of the first order up to the boundary  $\partial\Omega_\zeta$ , then the following analogues of the Gauss–Ostrogradsky formula is true:

$$\begin{aligned}
 &\int_{\partial\Omega_\zeta} \varphi(\zeta) \sigma \psi(\zeta) \\
 &= \int_{\Omega_\zeta} \left( \frac{\partial\varphi}{\partial x} \psi + \varphi \frac{\partial\psi}{\partial x} + \frac{\partial\varphi}{\partial y} i_2 \psi + \varphi i_3 \frac{\partial\psi}{\partial y} + \frac{\partial\varphi}{\partial z} i_3 \psi + \varphi i_3 \frac{\partial\psi}{\partial z} \right) dx dy dz.
 \end{aligned} \tag{3.3}$$

Using the equality (3.3) and the conditions (2.2), (2.3) we obtain the following theorem.

**Theorem 3.1.** *Suppose that a domain  $\Omega_\zeta$  has a closed piece-smooth boundary  $\partial\Omega_\zeta$  and  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is a right- $G$ -monogenic mapping in  $\Omega_\zeta$ , and  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is a left- $G$ -monogenic in  $\Omega_\zeta$  and continuous together with partial derivatives of the first order up to the boundary*

$\partial\Omega_\zeta$ . Then

$$\begin{aligned} & \int_{\partial\Omega_\zeta} \widehat{\Phi}(\zeta) \sigma \Phi(\zeta) \\ &= \int_{\Omega_\zeta} \left[ \widehat{\Phi}'(\zeta)(1 + i_2^2 + i_3^2)\Phi(\zeta) + \widehat{\Phi}(\zeta)(1 + i_2^2 + i_3^2)\Phi'(\zeta) \right] dx dy dz. \quad (3.4) \end{aligned}$$

*Proof.* Using the conditions (2.2), (2.3) we have

$$\begin{aligned} & \int_{\partial\Omega_\zeta} \widehat{\Phi}(\zeta) \sigma \Phi(\zeta) \\ &= \int_{\Omega_\zeta} \left( \widehat{\Phi}' \Phi + \widehat{\Phi} \Phi' + \widehat{\Phi}' i_2^2 \Phi + \widehat{\Phi} i_2^2 \Phi' + \widehat{\Phi}' i_3^2 \Phi + \widehat{\Phi} i_3^2 \Phi' \right) dx dy dz \\ &= \int_{\Omega_\zeta} \left[ (\widehat{\Phi}' + \widehat{\Phi}' i_2^2 + \widehat{\Phi}' i_3^2) \Phi + \widehat{\Phi} (\Phi' + i_2^2 \Phi' + i_3^2 \Phi') \right] dx dy dz \\ &= \int_{\Omega_\zeta} \left[ \widehat{\Phi}' (1 + i_2^2 + i_3^2) \Phi + \widehat{\Phi} (1 + i_2^2 + i_3^2) \Phi' \right] dx dy dz. \end{aligned}$$

□

The following statement is a consequence of Theorem 3.1.

**Theorem 3.2.** Under conditions of Theorem 3.1 with the additional assumption  $1 + i_2^2 + i_3^2 = 0$ , i. e. mappings  $\Phi$  and  $\widehat{\Phi}$  are solutions of the three-dimensional Laplace equation, the equality (3.4) can be rewritten in the form

$$\int_{\partial\Omega_\zeta} \widehat{\Phi}(\zeta) \sigma \Phi(\zeta) = 0.$$

#### 4. Cauchy integral theorem for a curvilinear integral

Let  $\gamma_\zeta$  be a Jordan rectifiable curve in  $E_3$ . For continuous mappings  $\varphi : \gamma_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and  $\psi : \gamma_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  of the forms (3.1) and (3.2), respectively, where  $(x, y, z) \in \gamma$ ,  $U_k : \gamma \rightarrow \mathbb{R}$ ,  $V_k : \gamma \rightarrow \mathbb{R}$  and  $P_m : \gamma \rightarrow \mathbb{R}$ ,  $Q_m : \gamma \rightarrow \mathbb{R}$ , we define a *curvilinear integral* along a Jordan rectifiable curve  $\gamma_\zeta$  by the equality:

$$\begin{aligned}
 \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) &:= \sum_{k,m=1}^4 e_k e_m \int_{\gamma} (U_k P_m - V_k Q_m) dx \\
 &+ \sum_{k,m=1}^4 e_k i_2 e_m \int_{\gamma} (U_k P_m - V_k Q_m) dy \\
 &+ \sum_{k,m=1}^4 e_k i_3 e_m \int_{\gamma} (U_k P_m - V_k Q_m) dz \\
 &+ i \sum_{k,m=1}^4 e_k e_m \int_{\gamma} (V_k P_m - U_k Q_m) dx \\
 &+ i \sum_{k,m=1}^4 e_k i_2 e_m \int_{\gamma} (V_k P_m - U_k Q_m) dy \\
 &+ i \sum_{k,m=1}^4 e_k i_3 e_m \int_{\gamma} (V_k P_m - U_k Q_m) dz,
 \end{aligned}$$

where  $d\zeta := dx + i_2 dy + i_3 dz$ .

If mappings  $\varphi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and  $\psi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  are continuous together with partial derivatives of the first order in a domain  $\Omega_\zeta$  and  $\Sigma_\zeta$  is an arbitrary piece-smooth surface in  $\Omega_\zeta$  with a rectifiable Jordan edge  $\gamma_\zeta$ , then the following analogue of *the Stokes formula* is true:

$$\begin{aligned}
 \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) &= \int_{\Sigma_\zeta} \left( \frac{\partial \varphi}{\partial x} i_2 \psi + \varphi i_2 \frac{\partial \psi}{\partial x} - \frac{\partial \varphi}{\partial y} \psi - \varphi \frac{\partial \psi}{\partial y} \right) dx dy \\
 &+ \left( \frac{\partial \varphi}{\partial y} i_3 \psi + \varphi i_3 \frac{\partial \psi}{\partial y} - \frac{\partial \varphi}{\partial z} i_2 \psi - \varphi i_2 \frac{\partial \psi}{\partial z} \right) dy dz \\
 &+ \left( \frac{\partial \varphi}{\partial z} \psi + \varphi \frac{\partial \psi}{\partial z} - \frac{\partial \varphi}{\partial x} i_3 \psi - \varphi i_3 \frac{\partial \psi}{\partial x} \right) dz dx. \tag{4.1}
 \end{aligned}$$

In the next theorem we show that the right-hand side of the equality (4.1) equals zero for the right- $G$ -monogenic mapping  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and the left- $G$ -monogenic mapping  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$ . Note that the following theorem is a generalization of Theorem 1 of [1].

**Theorem 4.1.** *Suppose that  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is a right- $G$ -monogenic mapping and  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is a left- $G$ -monogenic mapping in a domain  $\Omega_\zeta$ , and  $\gamma_\zeta$  is a rectifiable Jordan edge of some piece-smooth surface in  $\Omega_\zeta$ . Then*

$$\int_{\gamma_\zeta} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = 0. \tag{4.2}$$

*Proof.* Using the formula (4.1) and the conditions (2.2) and (2.3), we obtain

$$\begin{aligned} \int_{\gamma_\zeta} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) &= \int_{\Sigma_\zeta} \left( \frac{\partial \widehat{\Phi}}{\partial x} i_2 \Phi + \widehat{\Phi} i_2 \frac{\partial \Phi}{\partial x} - \frac{\partial \widehat{\Phi}}{\partial y} \Phi - \widehat{\Phi} \frac{\partial \Phi}{\partial y} \right) dx dy \\ &\quad + \left( \frac{\partial \widehat{\Phi}}{\partial y} i_3 \Phi + \widehat{\Phi} i_3 \frac{\partial \Phi}{\partial y} - \frac{\partial \widehat{\Phi}}{\partial z} i_2 \Phi - \widehat{\Phi} i_2 \frac{\partial \Phi}{\partial z} \right) dy dz \\ &\quad + \left( \frac{\partial \widehat{\Phi}}{\partial z} \Phi + \widehat{\Phi} \frac{\partial \Phi}{\partial z} - \frac{\partial \widehat{\Phi}}{\partial x} i_3 \Phi - \widehat{\Phi} i_3 \frac{\partial \Phi}{\partial x} \right) dz dx \\ &= \int_{\Sigma_\zeta} \left( \widehat{\Phi}'(\zeta) i_2 \Phi(\zeta) + \widehat{\Phi}(\zeta) i_2 \Phi'(\zeta) - \widehat{\Phi}'(\zeta) i_2 \Phi(\zeta) - \widehat{\Phi}(\zeta) i_2 \Phi'(\zeta) \right) dx dy \\ &\quad + \left( \widehat{\Phi}'(\zeta) i_2 i_3 \Phi(\zeta) + \widehat{\Phi}(\zeta) i_3 i_2 \Phi'(\zeta) - \widehat{\Phi}'(\zeta) i_3 i_2 \Phi(\zeta) - \widehat{\Phi}(\zeta) i_2 i_3 \Phi'(\zeta) \right) dy dz \\ &\quad + \left( \widehat{\Phi}'(\zeta) i_3 \Phi(\zeta) + \widehat{\Phi}(\zeta) i_3 \Phi'(\zeta) - \widehat{\Phi}'(\zeta) i_3 \Phi(\zeta) - \widehat{\Phi}(\zeta) i_3 \Phi'(\zeta) \right) dz dx = 0. \end{aligned}$$

□

We understand a triangle  $\Delta_\zeta$  as a plane figure bounded by three line segments connecting three its vertices. Denote by  $\partial\Delta_\zeta$  the boundary of the triangle  $\Delta_\zeta$  in the relative topology of its plane. Also we assume that the triangle  $\Delta_\zeta$  includes the boundary  $\partial\Delta_\zeta$ .

Since every triangle  $\Delta_\zeta \subset \Omega_\zeta$  can be included into a convex subset of a domain  $\Omega_\zeta$ , the following statement is a consequence of Theorem 4.1.

**Corollary 4.1.** *If  $\Omega_\zeta \subset E_3$  is a convex domain, a mapping  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is right- $G$ -monogenic and a mapping  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is left- $G$ -monogenic, then for an arbitrary triangle  $\Delta_\zeta$  such that  $\overline{\Delta_\zeta} \subset \Omega_\zeta$ , the following equality is true:*

$$\int_{\partial\Delta_\zeta} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = 0. \tag{4.3}$$

Let us consider the algebra  $\widetilde{\mathbb{H}}(\mathbb{R})$  with the basis  $\{e_k, ie_k\}_{k=1}^4$  over the field  $\mathbb{R}$  which is isomorphic to the algebra  $\mathbb{H}(\mathbb{C})$  over the field  $\mathbb{C}$ . In the algebra  $\widetilde{\mathbb{H}}(\mathbb{R})$  there exist another basis  $\{i_k\}_{k=1}^8$ , where the vectors  $i_1, i_2, i_3$  are the same as in the equalities (2.1).

For the element  $a := \sum_{k=1}^8 a_k i_k$ ,  $a_k \in \mathbb{R}$ , we define the Euclidian norm

$$\|a\| := \sqrt{\sum_{k=1}^8 a_k^2}.$$

Accordingly,  $\|\zeta\| = \sqrt{x^2 + y^2 + z^2}$  and  $\|i_1\| = \|i_2\| = \|i_3\| = 1$ .

Now we apply a scheme of the proof of the corresponding lemma for a function given in the complex plane (see, e. g., [7]) to the proof of the following statement.

**Lemma 4.1.** *Suppose that  $\varphi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and  $\psi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  are continuous mappings in a simply connected domain  $\Omega_\zeta$ , and  $\gamma_\zeta$  is a rectifiable curve in  $\Omega_\zeta$ . Then for an arbitrary  $\varepsilon > 0$  there exists a broken line  $\Lambda_\zeta \subset \Omega_\zeta$ , vertexes of which lie on the curve  $\gamma_\zeta$ , and such that*

$$\left\| \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - \int_{\Lambda_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) \right\| < \varepsilon. \tag{4.4}$$

*Proof.* Let us consider a closed domain  $\overline{D}_\zeta \subset \Omega_\zeta$ , containing inside the curve  $\gamma_\zeta$ . Since  $\varphi$  and  $\psi$  are continuous at every point of the domain  $\overline{D}_\zeta$ , then it is uniformly continuous in this domain. It means that the product of these mappings is uniformly continuous too. Thus, for an arbitrary  $\varepsilon_1 > 0$  there exists a number  $\delta(\varepsilon) > 0$  such that

$$\|\varphi(\zeta') \psi(\zeta') - \varphi(\zeta'') \psi(\zeta'')\| < \varepsilon_1, \tag{4.5}$$

if  $|\zeta' - \zeta''| < \delta(\varepsilon)$ , where  $\zeta', \zeta''$  are any points of the domain  $\overline{D}_\zeta$ . In addition, under the same assumptions, the following inequalities are true:

$$\|\varphi(\zeta') i_2 \psi(\zeta') - \varphi(\zeta'') i_2 \psi(\zeta'')\| < \varepsilon_2, \tag{4.6}$$

$$\|\varphi(\zeta') i_3 \psi(\zeta') - \varphi(\zeta'') i_3 \psi(\zeta'')\| < \varepsilon_3. \tag{4.7}$$

Let us divide the curve  $\gamma_\zeta$  into the  $n$  arcs  $Q_\zeta^0, Q_\zeta^1, \dots, Q_\zeta^{n-1}$  so that the length of each of them was less than  $\delta$  and enter the broken curve  $\Lambda_\zeta$  so that their broken links  $L_\zeta^0, L_\zeta^1, \dots, L_\zeta^{n-1}$  tied these arcs. By  $\zeta_0, \zeta_1, \dots, \zeta_{n-1}, \zeta_n$  denote the vertexes of the broken curve  $\Lambda_\zeta$ . Since the

length of every arc  $Q_\zeta^k$  is less than  $\delta$ , the distance between any two points on the same arc especially less than  $\delta$ . The same is true for links  $L_\zeta^k$ .

We compare the value of integral along the curve  $\gamma_\zeta$  with the value of the same integral along the broken curve  $\Lambda_\zeta$ . For this goal we consider a sum, which is an approximate value of the integral  $\int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta)$ :

$$S := \varphi(\zeta_0) \Delta\zeta_0 \psi(\zeta_0) + \varphi(\zeta_1) \Delta\zeta_1 \psi(\zeta_1) + \cdots + \varphi(\zeta_{n-1}) \Delta\zeta_{n-1} \psi(\zeta_{n-1}). \quad (4.8)$$

Since  $\Delta\zeta_k = \int_{Q_\zeta^k} d\zeta$ , the equality (4.8) can be represented in the form

$$S := \int_{Q_\zeta^0} \varphi(\zeta_0) d\zeta \psi(\zeta_0) + \int_{Q_\zeta^1} \varphi(\zeta_1) d\zeta \psi(\zeta_1) + \cdots + \int_{Q_\zeta^{n-1}} \varphi(\zeta_{n-1}) d\zeta \psi(\zeta_{n-1}). \quad (4.9)$$

On the other hand, the integral  $\int_{\gamma_\zeta} \varphi(\zeta) d\zeta \Psi(\zeta)$  can be represented in the form of the sum of the integrals along the arcs  $Q_\zeta^k$ :

$$\begin{aligned} \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) &= \int_{Q_\zeta^0} \varphi(\zeta) d\zeta \psi(\zeta) \\ &+ \int_{Q_\zeta^1} \varphi(\zeta) d\zeta \psi(\zeta) + \cdots + \int_{Q_\zeta^{n-1}} \varphi(\zeta) d\zeta \psi(\zeta). \end{aligned} \quad (4.10)$$

Consider the difference of the equations (4.10) and (4.9):

$$\begin{aligned} \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - S &= \int_{Q_\zeta^0} \left( \varphi(\zeta) d\zeta \psi(\zeta) - \varphi(\zeta_0) d\zeta \psi(\zeta_0) \right) \\ &+ \int_{Q_\zeta^1} \left( \varphi(\zeta) d\zeta \psi(\zeta) - \varphi(\zeta_1) d\zeta \psi(\zeta_1) \right) \\ &+ \cdots + \int_{Q_\zeta^{n-1}} \left( \varphi(\zeta) d\zeta \psi(\zeta) - \varphi(\zeta_{n-1}) d\zeta \psi(\zeta_{n-1}) \right) \\ &= \int_{Q^0} \left( \varphi(\zeta) \psi(\zeta) - \varphi(\zeta_0) \psi(\zeta_0) \right) dx + \int_{Q^0} \left( \varphi(\zeta) i_2 \psi(\zeta) - \varphi(\zeta_0) i_2 \psi(\zeta_0) \right) dy \end{aligned}$$

$$\begin{aligned}
& + \int_{Q^0} \left( \varphi(\zeta) i_3 \psi(\zeta) - \varphi(\zeta_0) i_3 \psi(\zeta_0) \right) dz \\
& + \dots + \int_{Q^{n-1}} \left( \varphi(\zeta) \psi(\zeta) - \varphi(\zeta_0) \psi(\zeta_0) \right) dx \\
& + \int_{Q^{n-1}} \left( \varphi(\zeta) i_2 \psi(\zeta) - \varphi(\zeta_0) i_2 \psi(\zeta_0) \right) dy \\
& + \int_{Q^{n-1}} \left( \varphi(\zeta) i_3 \psi(\zeta) - \varphi(\zeta_0) i_3 \psi(\zeta_0) \right) dz.
\end{aligned}$$

Since on the every arc  $Q_\zeta^k$  the inequalities (4.5) – (4.7) are true, we obtain

$$\left\| \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - S \right\| < (\varepsilon_1 Q_x^0 + \varepsilon_2 Q_y^0 + \varepsilon_3 Q_z^0) + \dots$$

$$\dots + (\varepsilon_1 \cdot Q_x^{n-1} + \varepsilon_2 Q_y^{n-1} + \varepsilon_3 Q_z^{n-1}) < \varepsilon Q^0 + \dots + \varepsilon Q^{n-1} < \varepsilon L, \quad (4.11)$$

where  $Q_x^j, Q_y^j, Q_z^j$  are lengths of the projections of the arc  $Q^j$  into the axes  $Ox, Oy, Oz$ , respectively,  $\varepsilon := \max\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$  and  $L$  is the length of the curve  $\gamma_\zeta$ .

In the same way we estimate the difference  $\int_{\Lambda_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - S$  and obtain

$$\left\| \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - S \right\| < \varepsilon L. \quad (4.12)$$

Taking into account the inequalities (4.11) and (4.12), we have

$$\begin{aligned}
\left\| \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - \int_{\Lambda_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) \right\| & \leq \left\| \int_{\gamma_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) - S \right\| \\
& + \left\| S - \int_{\Lambda_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) \right\| < 2\varepsilon L.
\end{aligned}$$

□

Now, using Corollary 4.1 and Lemma 4.1, we prove the following analogue of the Cauchy theorem for an arbitrary rectifiable curve in a convex domain.

**Theorem 4.2.** *Suppose that  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is a right- $G$ -monogenic mapping and  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  is a left- $G$ -monogenic mapping in a convex domain  $\Omega_\zeta$ . Then for any closed rectifiable Jordan curve  $\gamma_\zeta \subset \Omega_\zeta$  the equality (4.2) is true.*

*Proof.* **Basing** on Lemma 4.1, we inscribe the broken curve  $\Lambda_\zeta$  into the curve  $\gamma_\zeta$  so that the inequality (4.4) holds. Then we divide  $\Lambda_\zeta$  into triangles by the diagonals starting from a fixed vertex of  $\Lambda_\zeta$ . Since the domain  $\Omega_\zeta$  is convex, all obtained triangles contained in  $\Omega_\zeta$ . By Corollary 4.1, the integral along the every triangle equals to zero. Then the integral along the broken curve equals to zero too:

$$\int_{\Lambda_\zeta} \varphi(\zeta) d\zeta \psi(\zeta) = 0. \tag{4.13}$$

Now, the equality (4.2) is a consequence of the relations (4.4) and (4.13). □

In the case where  $\Omega_\zeta$  is an arbitrary domain, similarly to the proof of Theorem 3.2 [8], we can prove the following statement.

**Theorem 4.3.** *Let  $\Phi : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  be a right- $G$ -monogenic mapping and  $\widehat{\Phi} : \Omega_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  be a left- $G$ -monogenic mapping in a domain  $\Omega_\zeta$ . Then for every closed Jordan rectifiable curve  $\gamma_\zeta$  homotopic to a point in  $\Omega_\zeta$ , the equality (4.2) is true.*

*Proof.* Let a curve  $\gamma_\zeta$  be defined by the equality  $\zeta = \phi(t)$ ,  $0 \leq t \leq 1$ , where  $\phi(0) = \phi(1) = \zeta_0$ , and let  $\gamma_\zeta$  be homotopic to the point  $\zeta_0$ . Then there exists a continuous on the square  $Q := [0, 1] \times [0, 1]$  mapping  $H(s, t)$  of two real variables  $s$  and  $t$ , which takes values in the domain  $\Omega_\zeta$  and such that

$$H(0, t) = \phi(t), \quad H(1, t) \equiv \zeta_0 \quad \forall t \in [0, 1],$$

$$H(s, 0) = H(s, 1) = \zeta_0 \quad \forall s \in [0, 1].$$

Since the mapping  $H$  is continuous on a compact set  $Q$ , its image  $K := \{H(s, t) : (s, t) \in Q\}$  is a compact set in  $\Omega_\zeta$ .

Denote by  $\rho := \min_{\zeta' \in K, \zeta'' \in \partial\Omega_\zeta} \|\zeta' - \zeta''\|$ .

The mapping  $H$  is also uniformly continuous on the set  $Q$ . It means that there exists  $\delta > 0$  such that

$$\forall (s, t), (s', t') : |s' - s| < \delta, |t' - t| < \delta \Rightarrow \|H(s', t') - H(s, t)\| < \frac{\rho}{2}. \tag{4.14}$$

Let us choose a set of numbers  $0 = t_0 < t_1 < \dots < t_n = 1$ , which are satisfying the inequalities  $t_j - t_{j-1} < \delta$ ,  $j = 1, 2, \dots, n$ , and put  $s_1 = t_1$ . Let  $\zeta_{0,j} := H(0, t_j)$ ,  $\zeta_{1,j} := H(s_1, t_j)$  for  $j = 1, 2, \dots, n - 1$  and denote by  $L_\zeta^j$  a segment, beginning at the point  $\zeta_{0,j}$  and ending at the point  $\zeta_{1,j}$ . Also consider a curve  $\gamma_\zeta^{[1]} := \{H(s_1, t) : 0 \leq t \leq 1\}$ .

For a Jordan oriented curve  $\gamma_\zeta$ , by  $\gamma_\zeta[\zeta_1, \zeta_2]$  denote the arc beginning at the point  $\zeta_1$  and ending at the point  $\zeta_2$ .

Since of the inequality (4.14), the arcs  $\gamma_\zeta[\zeta_0, \zeta_{01}]$ ,  $\gamma_\zeta^{[1]}[\zeta_0, \zeta_{11}]$  and the segment  $L_\zeta^1$  are contained in the ball  $S(\zeta_0) := \{\zeta \in E_3 : \|\zeta - \zeta_0\| < \rho\}$ . Since  $S(\zeta_0)$  is a convex set and is contained in the domain  $\Omega_\zeta$ , the following equality is a consequence of Theorem 4.2

$$\int_{\gamma_\zeta[\zeta_0, \zeta_{01}]} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) + \int_{L_\zeta^1} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \int_{\gamma_\zeta^{[1]}[\zeta_0, \zeta_{11}]} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta). \quad (4.15)$$

The next inequalities follows from the inequalities (4.14):

$$\|\zeta - \zeta_{0,j}\| < \frac{\rho}{2} \quad \forall \zeta \in \gamma_\zeta[\zeta_{0,j}, \zeta_{0,j+1}],$$

$$\|\zeta - \zeta_{1,j}\| < \frac{\rho}{2} \quad \forall \zeta \in \gamma_\zeta^{[1]}[\zeta_{1,j}, \zeta_{1,j+1}], \quad \|\zeta_{1,j} - \zeta_{0,j}\| < \frac{\rho}{2}$$

for  $j = 1, 2, \dots, n - 2$ . Then the arcs  $\gamma_\zeta[\zeta_{0,j}, \zeta_{0,j+1}]$ ,  $\gamma_\zeta^{[1]}[\zeta_{1,j}, \zeta_{1,j+1}]$  and the segments  $L_\zeta^j, L_\zeta^{j+1}$  are contained in the ball  $S(\zeta_{0,j}) := \{\zeta \in E_3 : \|\zeta - \zeta_{0,j}\| < \rho\}$  for  $j = 1, 2, \dots, n - 2$ . Since  $S(\zeta_{0,j})$  is a convex set and is contained in  $\Omega_\zeta$ , the next equalities follows from the Theorem 4.2

$$\begin{aligned} & - \int_{L_\zeta^j} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) + \int_{\gamma_\zeta[\zeta_{0,j}, \zeta_{0,j+1}]} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) + \\ & + \int_{L_\zeta^{j+1}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \int_{\gamma_\zeta^{[1]}[\zeta_{1,j}, \zeta_{1,j+1}]} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) \end{aligned} \quad (4.16)$$

for  $j = 1, 2, \dots, n - 2$ .

Finally, similarly to the equality (4.15) we obtain the equality

$$- \int_{L_\zeta^{n-1}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) + \int_{\gamma_\zeta[\zeta_{0,n-1}, \zeta_0]} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \int_{\gamma_\zeta^{[1]}[\zeta_{1,n-1}, \zeta_0]} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta). \quad (4.17)$$

Adding all the equalities (4.15)–(4.17), we obtain the equality

$$\int_{\gamma_\zeta} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \int_{\gamma_\zeta^{[1]}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) \quad (4.18)$$

Then we put  $s_j = t_j$  and consider the curve  $\gamma_\zeta^{[j]} := \{H(s_j, t) : 0 \leq t \leq 1\}$  for  $j = 1, 2, \dots, n$ . Similarly to the equality (4.18), we obtain the equalities

$$\int_{\gamma_\zeta^{[1]}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \int_{\gamma_\zeta^{[2]}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \dots = \int_{\gamma_\zeta^{[n]}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta).$$

Hence, we have

$$\int_{\gamma_\zeta} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = \int_{\gamma_\zeta^{[n]}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta),$$

where the curve  $\gamma_\zeta^{[n]}$  degenerates to the point, because  $H(1, t) \equiv \zeta_0$ . Now, taking into account the equality

$$\int_{\gamma_\zeta^{[n]}} \widehat{\Phi}(\zeta) d\zeta \Phi(\zeta) = 0,$$

we complete the proof of the theorem.  $\square$

Now, let us consider a curvilinear Cauchy integral theorem for  $G$ -monogenic mappings in the case where a curve of integration lies on the boundary of a domain of  $G$ -monogeneity.

Let a closed Jordan rectifiable curve  $\gamma_\zeta \equiv \gamma_\zeta(t)$ , where  $0 \leq t \leq 1$ , which is homotopic to an interior point  $\zeta_0 \in \Omega_\zeta$ , be given on the boundary  $\partial\Omega_\zeta$  of the domain  $\Omega_\zeta$ . It means that there exists a mapping  $H(s, t)$ , which is continuous on the square  $[0, 1] \times [0, 1]$ , and such that  $H(0, t) = \gamma_\zeta(t)$ ,  $H(1, t) \equiv \zeta_0$ , and all curves  $\gamma_\zeta^s \equiv \gamma_\zeta^s(t) := \{\zeta = H(s, t) : 0 \leq t \leq 1\}$  for  $0 < s < 1$  are contained in the domain  $\Omega_\zeta$ .

Consider also the curves  $\Gamma_\zeta^t \equiv \Gamma_\zeta^t(s) := \{\zeta = H(s, t) : 0 \leq s \leq 1\}$ . By  $m$  denote the linear Lebesgue measure of a rectifiable curve.

The following theorem can be proved similarly to the proof of Theorem 2 in [2] and Theorem 4 in [9].

**Theorem 4.4.** *Suppose that  $\Phi : \bar{\Omega}_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  and  $\widehat{\Phi} : \bar{\Omega}_\zeta \rightarrow \mathbb{H}(\mathbb{C})$  are continuous mapping in the closure  $\bar{\Omega}_\zeta$  of a domain  $\Omega_\zeta$ ,  $\Phi$  is right- $G$ -monogenic and  $\widehat{\Phi}$  is left- $G$ -monogenic mapping in  $\Omega_\zeta$ . Suppose also that  $\gamma_\zeta \subset \partial\Omega_\zeta$  is any closed Jordan rectifiable curve homotopic to a point  $\zeta_0 \in \Omega_\zeta$  such that the curves of the family  $\{\Gamma_\zeta^t : 0 \leq t \leq 1\}$  are rectifiable and the set  $\{\text{mes} \gamma_\zeta^s : 0 \leq s \leq 1\}$  is bounded. Then the equality (4.2) is true.*

## References

- [1] V. S. Shpakivskyi, T. S. Kuzmenko, *Integral theorems for the quaternionic  $G$ -monogenic mappings* // An. Șt. Univ. Ovidius Constanța, **24** (2) (2016), 271–281.
- [2] T. S. Kuzmenko, *Curvilinear integral theorem for  $G$ -monogenic mappings in the algebra of complex quaternion* // Int. J. Adv. Research Math., **6** (2016), 21–25.
- [3] V. S. Shpakivskyi, T. S. Kuzmenko, *On monogenic mappings of a quaternionic variable* // J. Math. Sci., **221** (5) (2017), 712–726.
- [4] O. F. Herus, *On hyperholomorphic functions of the space variable* // Ukr. Math. J., **63** (4) (2011), 530–537.
- [5] V. V. Kravchenko, M. V. Shapiro, *Integral representations for spatial models of mathematical physics*, Pitman Research Notes in Mathematics, Addison Wesley Longman Inc, 1996.
- [6] V. S. Shpakivskyi, T. S. Kuzmenko, *On one class of quaternionic mappings* // Ukr. Math. J., **68** (1) (2016), 127–143.
- [7] I. I. Privalov, *Introduction to the Theory of Functions of a Complex Variable*, GITTL, Moscow, 1977. [in Russian]
- [8] E. K. Blum, *A theory of analytic functions in banach algebras* // Trans. Amer. Math. Soc., **78** (1955), 343–370.
- [9] S. A. Plaksa, V. S. Shpakivskyi, *Integral theorems for differentiable functions in a threedimensional harmonic algebra* // Reports of the NAS of Ukraine, **5** (2010), 23–30 [in Russian].

## CONTACT INFORMATION

**Tetyana Sergiivna Kuzmenko** Institute of Mathematics  
of the NAS of Ukraine,  
Kiev, Ukraine  
*E-Mail:* kuzmenko.ts15@gmail.com

**Vitalii Stanislavovich Shpakivskyi** Institute of Mathematics  
of the NAS of Ukraine,  
Kiev, Ukraine  
*E-Mail:* shpakivskyi86@gmail.com