

. . , . . , . . ,  
 ,  
 .  
 ,  
 .  
 ,  
 ,  
 ,  
 : , *V. viviparus*, , ,  
 .

Obtained in the analysis of conchial variability in the framework of this study are of fundamental importance. In the investigated species there is a conchological difference between individuals of different sexes, and the nature and extent of these differences are quite different. In *Viviparus viviparus*, the differences between females and males relate, first of all, to the proportions of the turtle and, to a lesser extent, to absolute measurements. With the reliability of discrimination of individuals of different sexes is at a rather low level due to the low discrimination of males, which are more like females than females on males.

**Keywords:** variability, *V. viviparus*, female, male, shell, discriminant analysis.

. ,  
 ,  
 ( )  
 , , .  
 ,  
 , ,  
 . , — ,  
 . , ,  
 ,  
 ,  
 ,  
 ,

[2, 4].

[1] *V. viviparus*

*V. subpurpureus*

[3]

Viviparus ,  
*V. viviparus.*  
1920-

[1],

10

[1],

100%.

1

	%		
	96,3	526	20
	19,9	145	36
	77,3	671	56

1. *Vivipara fasciata* Mull / . . . // . – 1928. – 3. – 88 .
2. Kotrla B. Sexual dimorphism of shell shape and growth of *Villosa villosa* (Wright) and *Elliptio laterina* (Conrad) (Pelecypoda: Unionidae / B. Kotrla, F. C. James // J. Moll. Stud. – 1987. – 53. – P.13–23.
3. Minton R. L. Evidence of sexual shape dimorphism in *Viviparus* (Gastropoda: Viviparidae) / R. L. Minton, L. L. Wang // J. Mollusc. Stud. – 2011. – P.315-317.
4. Son M. H. Sexual dimorphism of *Nucella lapillus* (Gastropoda: Muricidae) in North Wales, UK / M. H. Son, R. N. Hughes // J. Mollusc Stud. – 2000. – V. 66. – P. 489–498.