

# SYNTHESIS A NEW 5-HYDROXYSUBSTITUTED TRIAZOLO[1,4]DIAZEPINES


Dorokhov V. I.<sup>1</sup>, Kemskii S.V.<sup>2</sup>, Bol'but, A.V.<sup>2</sup>, Vovk M. V.<sup>2</sup>

<sup>1</sup>Polissia National University, [vdorokhov2@gmail.com](mailto:vdorokhov2@gmail.com)

<sup>2</sup>Institute of Organic Chemistry, National Academy of Sciences of Ukraine

Azoloannelized [1,4]diazepines are synthetically and biologically interesting condensed heterocyclic systems [1,2]. The pharmacological significance of pyrazolo[1,4]diazepines is demonstrated by their use as active substances of the anxiolytic drug zolasepam [3–5] and the antidepressant zomethapine [6,7]. In addition, a number of pyrazolo[1,4]diazepines revealed selective phosphodiesterase inhibitors [8] and oxytocin receptor antagonists [9].

The method proposed earlier by us [10] and other authors [11] for the formation of the hydroxydiazepine cycle based on 5-amino-N-(2,2-dialkoxyethyl)-1H-pyrazole(imidazole)-4-carboxamides turned out to be acceptable for the production of new derivatives triazolo[1,4]diazepines. The key substrates for this reaction were the anionic cyclization of azides **1a-l** synthesized by us with N-(2,2-dimethoxyethyl)-2-cyanacetamide (**2**) 5-amino-N-(2,2-dimethoxyethyl)-1H-1,2,3-triazole-4-carboxamides **3a-l** (Scheme 1).



The method is general in nature and allows involving alkyl-, aryl- and heterylazides in the process of formation of the triazole cycle and under mild conditions in the presence of *t*-BuOK as a base leads to **3a-l** compounds with yields of 79-94% (Table 1). The N-functionally substituted aminotriazolcarboxamides **3a-l** in formic acid at room temperature undergo light intramolecular cyclization at room temperature to form 5-hydroxy substituted triazolo[4,5-*e*][1,4]diazepines **4a-l** with close to quantitative yields (Scheme 1, Table 1).

**Table 1**

Product yields **3, 4 a-l**

Azide	R	Product (yield, %)	Product (yield, %)
1a	Me <sub>2</sub> CHCH <sub>2</sub>	3a (79)	4a (81)
1b	PhCH <sub>2</sub>	3b (85)	4b (89)
1c	4-MeOC <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	3c (89)	4c (87)
1d	3-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>	3d (84)	4d (88)
1e	Ph	3e (81)	4e (92)
1f	4-ClC <sub>6</sub> H <sub>4</sub>	3f (93)	4f (95)
1g	4-MeC <sub>6</sub> H <sub>4</sub>	3g (88)	4g (91)
1h	4-MeOC <sub>6</sub> H <sub>4</sub>	3h (86)	4h (93)
1i	4-O <sub>2</sub> NC <sub>6</sub> H <sub>4</sub>	3i (91)	4i (96)
1j	2,4-F <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	3j (91)	4j (90)
1k	2,4-Me <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	3k (81)	4k (89)
1l	1-Methylpyrazole-3-yl	3l (92)	4l (94)

The IR spectra of 4a–l compounds are characterized by a set of absorption bands corresponding to the valence oscillations of the bonds C=O (1627–1636 cm<sup>-1</sup>), N–H (3247–3312 cm<sup>-1</sup>) and OH (3247–3312 cm<sup>-1</sup>). In the spectra of NMR 1H, along with typical signals of R substituents, there are multiplets of protons 6-CH<sub>2</sub> at 2.98–3.38 m. d., protons of 5-CH at 4.86–5.18 m. d., as well as signals of protons NH: doublets at 7.26–8.01 m. d. and multiplets at 7.13–7.66 m. d.

Thus, we show that intramolecular cyclization of 5-amino-N-(2,2-dimethoxyethyl)-1H-1,2,3-triazole-4-carboxamides in formic acid is a convenient way to synthesize 5-hydroxy[1,2,3]triazolo[4,5-e][1,4] diazepines, which are promising objects for synthetic and medicinal chemistry.

1. Ramajayam, R.; Girdhar, R.; Yadav, M. R. *Mini-Rev. Med.Chem.* **2007**, *7*, 793.
2. Кемский, С. В.; Ботьбуь, А. В.; Дмьтрьв, Ю. В.; Вовк, М. В. *Журн. орган. фарм. хьмьї* **2017**, *15*, 3.
3. DeWald, H. A.; Lobbestael, S.; Butler, D. E. *J. Med. Chem.* **1977**, *20*, 1562.
4. Sleeman, J. M.; Cameron, K.; Mudakikwa, A. B.; Nizeyi, J.-B.; Anderson, S.; Cooper, J. E.; Richardson, H. M.; Macfie, E. J.; Hastings, B.; Foster, J. W. *J. Zoo Wild. Med.* **2000**, *31*, 9.
5. Cattet, M. R.; Caulkett, N. A.; Polischuk, S. C.; Ramsay, M. A. *J. Zoo Wild. Med.* **1999**, *30*, 354.
6. Katz, R. J. *Pharmacol., Biochem. Behav.* **1984**, *21*, 487.
7. Fairchild, C. J.; Rush, A. J.; Vasavada, N.; Giles, D. E.; Khatami, M. *Psychiatry Res.* **1986**, *18*, 217.
8. Henriksson, K.; Lisius, A.; Sjö, P.; Storm, P. WO Patent 2007040435.
9. Hudson, P.; Pitt, G. R. W.; Batt, A. R.; Roe, M. B. US Patent 20070197608.
10. Bol'but, A. V.; Kemskii, S. V.; Vovk, M. V. *Russ. J. Org. Chem.* **2012**, *48*, 991. [*Журн. орган. хьмьї* **2012**, *48*, 994.]
11. Tantravedi, S.; Chakraborty, S.; Shah, N. H.; Fishbein, J. C.; Hosmane, R. S. *Bioorg. Med. Chem.* **2013**, *21*, 4893.