

On analogue of Koebe-Bloch theorem for ring homeomorphisms

Denys Romash

(Zhytomyr Ivan Franko State University)

E-mail: dromash@num8erz.eu

Evgeny Sevost'yanov

(Zhytomyr Ivan Franko State University; Institute of Applied Mathematics and Mechanics,
Slov'yans'k)

E-mail: esevostyanov2009@gmail.com

All definitions and notions used below may be found in [1]. Let $S_i = S(x_0, r_i) = \{x \in \mathbb{R}^n : |x - x_0| = r_i\}$, $i = 1, 2$, $n \geq 2$, and let $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ be a Lebesgue measurable function satisfying the condition $Q(x) \equiv 0$ for $x \in \mathbb{R}^n \setminus D$. In what follows, $M(\cdot)$ denotes the modulus of family of paths, and $dm(x)$ is an element of the Lebesgue measure in \mathbb{R}^n . A mapping $f : D \rightarrow \overline{\mathbb{R}^n}$, $\overline{\mathbb{R}^n} := \mathbb{R}^n \cup \{\infty\}$, is called a *ring Q -mapping at the point $x_0 \in \overline{D} \setminus \{\infty\}$* , if the condition

$$M(f(\Gamma(S_1, S_2, D))) \leq \int_{A \cap D} Q(x) \cdot \eta^n(|x - x_0|) dm(x),$$

$A = A(x_0, r_1, r_2) = \{x \in \mathbb{R}^n : r_1 < |x - x_0| < r_2\}$, holds for all $0 < r_1 < r_2 < d_0 := \sup_{x \in D} |x - x_0|$ and all Lebesgue measurable functions $\eta : (r_1, r_2) \rightarrow [0, \infty]$ such that

$$\int_{r_1}^{r_2} \eta(r) dr \geq 1.$$

Below $h(x, y)$ denotes the chordal (spherical metric) between points $x, y \in \overline{\mathbb{R}^n}$. Let $h(E)$ be achordal diameter of the set E in $\overline{\mathbb{R}^n}$. Given $x_0 \in \overline{\mathbb{R}^n}$ and $r > 0$ we set $B_h(x_0, r) := \{x \in \overline{\mathbb{R}^n} : h(x, x_0) < r\}$. For $x_0 \in \mathbb{R}^n$ and $r_0 > 0$, as usual, $B(x_0, r_0) = \{x \in \mathbb{R}^n : |x - x_0| < r_0\}$. A family \mathfrak{F} of mappings $f : D \rightarrow \overline{\mathbb{R}^n}$ is called *uniformly open at a point $x_0 \in D$* , if for every $\varepsilon_0 > 0$, $\varepsilon_0 < \text{dist}(x_0, \partial D)$, there is $r_0 = r_0(x_0, \varepsilon_0) > 0$ such that $B_h(f(x_0), r_0) \subset f(B(x_0, \varepsilon_0))$ for every $f \in \mathfrak{F}$. A family \mathfrak{F} of mappings $f : D \rightarrow \overline{\mathbb{R}^n}$ is called *uniformly open on K* , if for every $\varepsilon_0 > 0$ there exists $r_0 = r_0(K, \varepsilon_0) > 0$ such that $B_h(f(x_0), r_0) \subset f(B(x_0, \varepsilon_0))$ for every $f \in \mathfrak{F}$ and every $B(x_0, \varepsilon_0) \subset K$.

Given a domain D in \mathbb{R}^n , $n \geq 2$, a Lebesgue measurable function $Q : D \rightarrow [0, \infty]$, a compact set $K \subset D$ and a number $\delta > 0$ denote by $\mathfrak{F}_{K, Q}^\delta(D)$ a family of all ring Q -homeomorphisms $f : D \rightarrow \overline{\mathbb{R}^n}$ such that $h(f(K), \partial f(D)) = \inf_{x \in f(K), y \in \partial f(D)} h(x, y) \geq \delta$. The following result holds.

Theorem 1. (An analogue of Koebe-Bloch theorem). *Let D be a bounded domain. Assume that, for each point $x_0 \in D$ and for every $0 < r_1 < r_2 < r_0 := \sup_{x \in D} |x - x_0|$ there is a set $E_1 \subset [r_1, r_2]$ of a positive linear Lebesgue measure such that the function Q is integrable with respect to \mathcal{H}^{n-1} over the spheres $S(x_0, r)$ for every $r \in E_1$. Then the family $\mathfrak{F}_{K, Q}^\delta(D)$ is uniformly open on K .*

Remark 2. If f is a homeomorphism, $f : D \rightarrow \mathbb{R}^n$, $n \geq 3$, $f \in W_{\text{loc}}^{1, \varphi}$, $\int_1^\infty \left[\frac{t}{\varphi(t)} \right]^{\frac{1}{n-2}} dt < \infty$ and $K_O(x, f) \in L_{\text{loc}}^{n-1}$, then $g := f^{-1} \in W_{\text{loc}}^{1, n}$, where $K_O(x, f)$ is an outer dilatation of f at x . On

the other hand, if $K_I(y, g) \in L_{\text{loc}}^1$, then g is a ring Q_* -homeomorphism with $Q_* = K_I(y, g)$ (see Corollary 8.5, Theorem 8.6 in [1]). Thus, Theorem 1 holds for Orlicz-Sobolev classes $W_{\text{loc}}^{1,\varphi}$ under above assumptions.

REFERENCES

- [1] Martio O., Ryazanov V., Srebro U. and Yakubov E. *Moduli in Modern Mapping Theory*. Springer Science + Business Media, LLC : New York, 2009.

Acknowledgements. This talk is supported by the National Research Foundation of Ukraine (Project “Analogues of Carathéodory and Koebe-Bloch theorems for Orlicz-Sobolev classes”, Project number 2025.02/0010).